

**U. S. LOTIC PROPER FUNCTIONING CONDITION (PFC) CHECKLIST**  
**USER MANUAL**  
**(Current as of 5/20/2010)**

This user manual is intended to accompany the Lotic Proper Functioning Condition (PFC) Standard Checklist Form for assessing proper functioning condition of riparian wetlands associated with systems having flowing water and (usually) a defined channel. This document serves to assist data collectors in answering each item on the form and as an aid to the database user in the interpretation of data presented. Another form entitled the Lentic Proper Functioning Condition (PFC) Standard Checklist Form, with a different set of user guidelines, is to be used for lentic (still water) wetlands.

**BACKGROUND INFORMATION**

**Flowing Water (Lotic) Wetlands vs. Still Water (Lentic) Wetlands**

Cowardin and others (1979) point out that no single, correct definition for wetlands exists, primarily due to the nearly unlimited variation in hydrology, soil, and vegetative types. Wetlands are lands transitional between aquatic (water) and terrestrial (upland) ecosystems. Windell and others (1986) state, “wetlands are part of a continuous landscape that grades from wet to dry. In many cases, it is not easy to determine precisely where they begin and where they end.”

In the semi-arid and arid portions of western North America, a useful distinction has been made between wetland types based on association with different aquatic ecosystems. Several authors have used *lotic* and *lentic* to separate wetlands associated with flowing water from those associated with still water. The following definitions represent a synthesis and refinement of terminology from Shaw and Fredine (1956), Stewart and Kantrud (1972), Boldt and others (1978), Cowardin and others (1979), American Fisheries Society (1980), Johnson and Carothers (1980), Cooperrider and others (1986), Windell and others (1986), Environmental Laboratory (1987), Kovalchik (1987), Federal Interagency Committee for Wetland Delineation (1989), Mitsch and Gosselink (1993), and Kent (1994).

*Lotic wetlands* are associated with rivers, streams, and drainage ways. Such wetlands contain a defined channel and floodplain. The channel is an open conduit, which periodically or continuously carries flowing water, dissolved and suspended material. Beaver ponds, seeps, springs, and wet meadows on the floodplain of, or associated with, a river or stream are part of the lotic wetland.

*Lentic wetlands* are associated with still water systems. These wetlands occur in basins and lack a defined channel and floodplain. Included are permanent (i.e., perennial) or intermittent bodies of water such as lakes, reservoirs, potholes, marshes, ponds, and stockpools. Other examples include fens, bogs, wet meadows, and seeps not associated with a defined channel.

**Functional vs. Jurisdictional Wetland Criteria**

Defining wetlands has become more difficult as greater economic stakes have increased the involvement of more politics and less science. A universally accepted wetland definition satisfactory to all users has not yet been developed because the definition depends on the objectives and the field of interest. However, scientists generally agree that wetlands are characterized by one or more of the following features: 1) *wetland hydrology*, the driving force creating all wetlands, 2) *hydric soils*, an indicator of the absence of oxygen, and 3) *hydrophytic vegetation*, an indicator reflecting wetland site conditions. The problem is how to define and obtain consensus on thresholds for these three criteria and various combinations of them.

Wetlands are not easily identified and delineated for jurisdictional purposes. Functional definitions have generally been difficult to apply to the regulation of wetland dredging or filling. Although the intent of legislation is to protect wetland functions, delineation of jurisdictional wetlands has relied largely on structural features or attributes. The hydrogeomorphic (HGM) approach being developed by the US Corps of Engineers is intended to focus more specifically on wetland functions.

The prevailing view among many wetland scientists is that *functional wetlands need* to meet only one of the three criteria as outlined by Cowardin and others (1979) (e.g., hydric soils, hydrophytic plants, and wetland hydrology). On the other hand, *jurisdictional wetlands need to* meet all three criteria, except in limited situations. Even though functional wetlands may not meet jurisdictional wetland requirements, they certainly perform wetland functions resulting from the greater amount of water that accumulates on or near the soil surface relative to the adjacent uplands. Examples include some woody draws

occupied by the *Fraxinus pennsylvanica/Prunus virginiana* (green ash/common chokecherry) habitat type and some floodplain sites occupied by the *Artemisia cana/Agropyron smithii* (silver sagebrush/western wheatgrass) habitat type or the *Pinus ponderosa/Cornus stolonifera* (ponderosa pine/red-osier dogwood) habitat type. Currently, many of these sites fail to meet jurisdictional wetland criteria. Nevertheless, these sites do provide important wetland functions and may warrant special managerial consideration. The current interpretation, at least in the western United States, is that not all functional wetlands are jurisdictional wetlands, but all jurisdictional wetlands are functional wetlands.

### **Polygon Delineation**


The lotic wetland inventory process incorporates data on a wide range of biological and physical categories. The basic unit of delineation within which this data is collected is called a ***polygon***. A polygon is the area upon which one set of data is collected. One inventory form is completed (i.e., one set of data is collected) for each polygon. One or more (usually several) polygons constitute a project. A lotic (riparian) polygon is an area adjacent to a waterway (stream or river). Polygons are delineated on 7.5-minute topographic (topo) maps by marking the upper and lower ends before observers go to the field. (The widths of most riparian wetland zones are unknown before the inventory and cannot be pre-marked.) On 7.5-minute topo maps, most polygons are usually drawn as a single line following the stream or river and are numbered sequentially proceeding downstream. It is important to clearly mark and number the polygons on the topo map. Polygons are numbered pre-field (in the office) with consecutive integers (1, 2, 3 . . .). In cases where field inspection shows a need to change the delineation or to subdivide pre-drawn polygons, additional polygons should be numbered using alpha- numerics (e.g., 1a, 1b, 2a, 2b, etc.). When delineated polygons are subsequently combined in the field, the combinations are to be identified by the hyphenated tags of both combined parts (e.g., 1-2, 2-3, etc.).

If aerial photos are available, advance (pre-field) polygon delineations may be based on vegetation differences, geologic features, or other observable characteristics. On larger systems with wide riparian areas, aerial photos may allow the pre-field delineation of multiple polygons away from the channel. In these cases, where polygons can be drawn as enclosed units (instead of just as a line), a minimum mapping unit of 5 or 10 acres (2 to 4 ha) should be used. The size of the minimum mapping unit should be based on factors such as management capabilities and the costs and capabilities of data collection.

In the field, observers are to verify (ground truth) the office-delineated polygon boundaries. If the pre-assigned numbers are used, be sure the inventoried polygons correspond exactly as drawn. Observers are allowed to move polygon boundaries, create new polygons, or consolidate polygons if the vegetation, geography, location of fences, or width of the wetland zone justify it. If polygon boundaries are changed, the changes must be clearly marked on the field copies of the 7.5-minute topographic maps. The original polygon numbers should be retained on the map for cross-reference. ***Polygons should not cross fences between areas with different management.***

Upper and lower polygon boundaries are placed at distinct locations such as fences, stream confluences, or stream meanders that can be recognized in the field. Polygons should not cross fences between areas with different management. In most cases, polygons are delineated one quarter to three quarters of a mile long. On smaller streams, polygons include the land on both sides of the stream. On large rivers, or if property ownership or access differs, polygons may include only one side of a stream.

The outer boundaries of riparian polygons are at the wetland vegetative outer edges. These boundaries are sometimes clearly defined by abrupt differences in the geography and/or vegetation, but proper determination often depends on experienced interpretation of more subtle differences. The area to be assessed includes any terraces dominated by facultative wetland and wetter plant species (Reed 1988), the active floodplain, and streambanks. Reference to Reed's list of plants found in wetlands should not be necessary to determine the area for evaluation. The evaluator should simply focus on that area which is obviously more lush, dense, or greener by virtue of proximity to the stream.



**Figure 2.** A schematic example of a typical riparian zone cross section showing near-channel landform features. *Note:* FAC (facultative), OBL (obligate), UPL (upland), etc. refer to categories of frequency a species is found in wetlands (Reed 1988).

The location of the inner (or streamside) polygon boundary must be known (at least approximately), even on polygons that span the stream. On most streams the area of the channel bottom is excluded from the polygon. (*Note: The whole channel width extends from right bankfull stage to left bankfull stage; however we need to include the lower banks in all polygons, therefore consider for exclusion ONLY the relatively flat and lowest area of the channel—the “bottom.”*) This allows data to be collected on the riparian area while excluding the aquatic zone, or open water, of the stream. The aquatic zone is the area covered by water and lacking persistent emergent vegetation. Persistent emergent vegetation consists of perennial wetland species that normally remain standing at least until the beginning of next growing season, e.g., *Typha* species (cattails), *Scirpus* species (bulrushes), *Carex* species (sedges), and other perennial graminoids.

In many systems, large portions of the channel bottom may become exposed due to seasonal irrigation use, hydroelectric generation, and natural seasonal changes such as are found in many prairie ecosystems. In these cases, especially the prairie streams, the channel bottom may have varying amounts of herbaceous vegetation, and the channel area is **included** in the polygon as area to be inventoried. Typically, these are the “pooled channel” stream type that has scour pools scattered along the length, interspersed with reaches of grass, bulrush, or sedge-covered channel bottom. If over half (>50%) the channel bottom area has a canopy cover of persistent vegetation cover (perennial species), taken over the entire length of the polygon as a whole, then it qualifies for inclusion within the inventoried polygon area. If you are in doubt whether to include the channel bottom in the polygon, then leave it out, but be sure to indicate this in the comment section. This is important so that future assessments of the polygon will be looking at the same area of land.

## PFC FORM ITEM CODES AND INSTRUCTIONS

### Polygon Data

The following are the codes and instructions for the individual data items on the form. All data items are to be recorded in the field unless otherwise noted. Numbering corresponds to that of items on the form. Do not use “—” and do not leave items blank. *NA* means the item is not applicable to a particular polygon. *NC* means data was not collected for that item in a particular polygon. Observers must write legibly and should limit use of abbreviations throughout to prevent confusion.

**Record ID No.** This is the unique identifier allocated to each polygon. This number will be assigned in the office when the form is entered into the database.

### Administrative Data

**A1.** Agency or organization collecting the data.

**A2.** Funding Agency/Organization.

**A3a.** BLM (Bureau of Land Management) State Office.

**A3b.** BLM Field Office/Field Station.

**A3c.** BLM Office Code (recorded in the office).

**A3d.** Is the polygon in an active BLM grazing allotment (recorded in the office)?

**A3e, f.** For BLM polygons, the BLM Office Code, whether the polygon is in an active BLM grazing allotment, and the Allotment Number is supplied by the BLM. These items are entered into the computer in the office; the computer then references a master list of Allotment ID's to complete the remaining Allotment information. Because some polygons incorporate more than one Allotment, space is provided to enter two sets of Allotment information. The master Allotment list is periodically updated by the BLM National Applied Resource Sciences Center to make needed corrections.

**A4.** USDI Fish and Wildlife Service Refuge name.

**A5.** Indian Reservation name.

**A6.** USDI National Park Service Park/National Historical Site name.

**A7.** USFS (Forest Service) National Forest name.

**A8.** Other location.

**A9.** Year the field work was done.

**A10.** Date of field work by day, month, and year.

**A11.** Names of all field data observers.

**Note:** Information for items **A12a-h** is found in the office; field evaluators need not complete these items.

**A12.** The several parts of these items identify various ways in which a data record may represent a resampling of a polygon that may have been inventoried again at some other time. The data in this record may have been collected on an area that coincides precisely with an area inventoried at another time and recorded as another record in the database. It may also represent the resampling of only a part of an area previously sampled. This would include the case where this polygon overlaps, but does not precisely and entirely coincide with one inventoried at another time. One other case is where more than one polygon inventoried one year coincides with a single polygon inventoried another year. All of these cases are represented in the database, and all have some value for monitoring purposes, in that they give some information on how the status on a site changes over time. ***This is done in the office with access to the database; field evaluators need not complete these items.***

**A12a.** Has any part of the area within this polygon been inventoried previously, or subsequently, as represented by any other data record in the database? Such other records would logically carry different dates.

**A12b.** Does the areal extent of this polygon exactly coincide with that of any other inventory represented in the database? In many cases, subsequent inventories only partially overlap spatially. The purpose of this question is to identify those records that can be compared as representing exactly the same ground area.

**A12c.** Does this record represent the latest data recorded for this site (polygon)?

**A12d.** If A12b is answered "Yes," then enter the record ID number(s) of any other previous or subsequent re-inventories (resampling) of this exact polygon for purposes of cross-reference.

**A12e.** Enter the years of any records recorded in item A12d as representing other inventories of this exact polygon.

**A12f.** Even though this polygon is not a re-inventory of the exact same area as any other polygon, does it share at least some common area with one or more polygons inventoried at another time?

**A12g.** Enter the years of any other inventories of polygons sharing common ground area with this one.

**A12h.** If A12f is answered “Yes,” then enter the record ID number(s) of any other polygon(s) sharing common ground area with this one.

**A13a.** Has a management change been implemented on this polygon?

**A13b.** If A13a is answered “Yes,” in what year was the management change implemented?

**A13c.** If A13a is answered “Yes,” describe the management change implemented.

**Location Data**

**B1.** State in which the field work was done (recorded in the office).

**B2.** County or municipal district in which the field work was done (recorded in the office).

**B3.** This field for allotment or range unit is intended for entities other than the BLM to use for grouping polygons by management unit. The BLM management units are grouped using the grazing allotment information in A3 above.


**B4a.** For lentic polygons the area is usually listed as a lake name, or other local designation that identifies the area where the inventory is conducted. If possible, use a name that is shown on the 7.5 minute topographic map.

**B4b.** Record the stream (if there is one) with which the inventoried lentic wetland is associated. Such association may be by inlet or outlet surface flow, or by general ground water (sub surface) connection.

**B4c, d.** Polygons are grouped together for management purposes. For example, all polygons around Henry’s Lake in the Idaho Falls Field Office could be identified as Group Name: Idaho Falls Field Office; Group Number: 1 (recorded in the office).

**B5.** Polygon number is a sequential identifier of the portion of the area assessed. This is referenced to the map delineations. Sequences normally progress clockwise.

**B6.** The Township, Range, Section, 1/4 section, and 1/4–1/4 section is the location of the midpoint of the polygon area. When reading this information as a legal description, the order is presented from smallest to largest unit. Below is a schematic showing the quarter sections, with the SE quarter divided into quarter-quarters.



**B7.** Elevation (feet or meters) of the polygon midpoint. Elevation is interpolated from the topographic map(s).

**B8a-e.** The US Geological Survey has divided the nation into successively smaller hydrologic units based on drainage basins. These units in the United States are subdivided into fourth levels, uniquely identified by a two-digit number for each level. This results in a eight-digit identifier for a drainage at the fourth level. Some regions have units defined to the fifth and sixth level (finer scales). (Data is entered in the office.)

**B9a.** Record the latitude and longitude of the polygon, along with the GPS projection and accuracy. Record the degrees, minutes, and seconds. *Note: All of North America is latitude = North, and longitude = West.*

**B9b.** Record any comments pertaining to the location of the polygon.

**B10.** Record the name(s) of the 7.5 minute quadrangle map(s) locating the polygon using precisely the name listed on the map sheet. Provision is made for listing two maps in case the polygon crosses between two maps.

## Selected Summary Data

**C1.** Wetland type is a categorical description of the predominant polygon character. Select from the following list of categories that may occur within a lotic system the one that best characterizes the majority of the polygon. Observers will **select only one category** as representative of the entire polygon. If significant amounts of other categories are present, indicate this in Vegetation Comments (item D17) or consider dividing the original polygon into two or more polygons.

---

### Category Description

---

**Perennial Stream.** A stream or stretch of stream that flows continuously for most of most years. Perennial streams are generally fed in part by springs or discharge from groundwater. Perennial streams are distinguished from larger rivers by size. Streams wider than 50 ft (15 m) are considered rivers for the purpose of this inventory (see below).

**Intermittent Stream.** A stream or stretch of stream which flows only at certain periods of the year when it receives water from springs, discharge from groundwater, or melting snow in mountainous areas. These streams generally flow continuously at least one month most years.

**Ephemeral Stream.** A stream or stretch of stream that flows in normal water years only in direct response to precipitation. In normal years, it receives no water from springs and no extended supply from melting snow or other surface source. Ephemeral streams are not in contact with groundwater and normally do not flow continuously for as long as one month. Not all ephemeral streams support riparian plant communities.

**Subterranean Stream.** A stream that flows underground for part of the stream reach. This occurs on systems composed of coarse textured, porous substrates. Surface flow may disappear and re-emerge farther downstream.

**Pooled Channel Stream.** An intermittent stream that has significant channel pools after surface flow ceases. Pools are generally at meander curves and are usually considerably deeper than the rest of the channel bottom. Water sources for the pools may be springs or contact with subsurface groundwater. This stream type is typical of fine textured sedimentary plains in semi-arid regions where headwater drainages lack the extended runoff of deep mountain snowpack. This stream type may not be apparent early in the season when flow is continuous.

**River.** Rivers are generally larger than streams. They flow year around, in years of normal precipitation and when significant amounts of water are not being diverted out of them. Those watercourses called rivers on topographic quads and/or those having bankfull channel widths greater than 50 ft (15 m) will be classified as rivers for the purpose of this inventory.

**Beaver Dams.** A system that is predominantly characterized by beaver dams that change the character of the system from a regular flowing channel to a “stepped” system of ponds where water is spread wide and flow velocity is apparent only at each dam outlet before it enters the next pond. Water is still flowing through the riparian system.

**Wet Meadow.** This type of wetland may occur in either running water (lotic) or in still water (lentic) systems. A lotic wet meadow has a defined channel or flowing surface water nearby, but is typically much wider than the riparian zone associated with the classes described above. This is often the result of the influence of lateral groundwater not associated with the stream flow. Lotic and lentic wet meadows may occur in proximity (e.g., when enough groundwater emerges to begin to flow from a mountain meadow, the system goes from lentic to lotic). Such communities are typically dominated by herbaceous hydrophytic vegetation that requires saturated soils near the surface, but tolerates no standing water for most of the year. This type of wetland typically occurs as the filled-in basin of old beaver ponds, lakes, and potholes.

**Spring/Seep.** Groundwater discharge areas. In general, springs have more flow than seeps. This wetland type may occur in a running water (lotic) or still water (lentic) system.

**Irrigation Canal.** Includes all types of canals and ditches associated with irrigation systems.

**Other.** Describe the water source (e.g., irrigation return flow, industrial discharge, etc.).

**Upland.** This designation is for those areas which are included in the inventoried polygon, but which do not support functional wetland vegetation communities. Such areas may be undisturbed inclusions of naturally occurring high ground or such disturbed high ground as roadways and other elevated sites of human activity.

---

**C2.** The size (acres/hectares) of polygons large enough to be drawn as enclosed units on 1:20,000 or 1:50,000 scale maps is determined in the office using a planimeter, dot grid, or GIS. For polygons too small to be accurately drawn as enclosed units on the maps, and which are represented by line segments on the map along the drainage bottom, polygon size is calculated using polygon length and average polygon width (items C5 and C7a).

**C3a-d.** Evaluators may be asked to survey some areas that have not been determined to be wetlands for the purpose of making such a determination. Other polygons include areas supporting non-wetland vegetation types. A “Yes” answer here indicates that no part of the polygon keys to a riparian habitat type or community type (HT/CT). Areas classified in item D15 as any vegetation type described in a riparian and/or wetland classification document for the region in which you are working

are counted as functional wetlands. Areas listed as UNCLASSIFIED WETLAND TYPE are also counted as functional wetlands. Other areas are counted as non-wetlands, or uplands. The functional wetland fraction of the polygon area is listed in item C3c in acres and as a percentage of the entire polygon area in item C3d.

**C4.** Some riparian areas do not contain an unvegetated, defined stream channel. In some cases, these polygons are in ephemeral systems which may flow infrequently, but which do support riparian plant communities. In other cases, these polygons may be associated with larger river systems that have wide floodplains where polygons may be delineated in areas not adjacent to the channel.

**C5.** Channel length—the length of channel contained within or adjacent to the polygon—is measured by scaling from the map. This data is considered accurate to the nearest 0.1 mile (0.16 km).

**C6.** In some cases, the polygon record is used to characterize, or represent, a larger portion of a stream system. The length represented by the polygon is given here. For example, a 0.5-mile polygon may be used to represent 4 miles of a stream. In the case, 0.5 is the channel length of the polygon (item C5), and 4 miles is entered in item C6.

**C7a.** Record average width of the polygon, which on smaller streams corresponds to the width of the riparian zone. To determine this width, subtract the width of the non-vegetated stream channel (item F9) from the distance between the two opposite riparian/upland boundaries. In the case of very wide systems where the polygon inventoried does not extend across the full width of the riparian zone (e.g., area with riparian vegetation communities lies outside the polygon), record the average width of the polygon inventoried and make note of the situation in the narrative comments.

**C7b.** Record the range of width (ft/m), narrowest to widest, of the riparian zone in the polygon.

### **LOTIC PFC STANDARD CHECKLIST**

For assistance and clarification in filling out and interpreting the checklist, refer to USDI BLM Technical Reference 1737-9 (Prichard and others 1993, revised 1995) and USDI BLM Technical Reference 1737-15 (Prichard and others 1998).

## LITERATURE CITED

- American Fisheries Society, Western Division. 1980. Position paper on management and protection of western riparian stream ecosystems. American Fisheries Society, Bethesda, Maryland, USA. 24 p.
- Boldt, Charles D., Daniel W. Uresk, and Keith E. Severson. 1978. Riparian woodlands in jeopardy on Northern High Plains. In: Strategies for protection and management of floodplain wetlands and other riparian ecosystems (R. R. Johnson and J. F. McCormick, Technical Coordinators). USDA Forest Service General Technical Report WO-12. Washington, DC, USA. pp. 184-189.
- Cooperrider, Allen Y., Raymond J. Boyd, and Hanson R. Stuart. 1986. Inventory and monitoring of wildlife habitat. USDI Bureau of Land Management, Denver Service Center, Denver Colorado, USA. 858 p.
- Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of wetlands and deep water habitats of the United States. USDI Fish and Wildlife Service, Office of Biological Services, Washington, DC, USA. Publication Number FWS/OBS-79/31. 107 p.
- Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Technical Report Y-87-1. US Army Engineer Waterways Experiment Station, Vicksburg, Mississippi, USA. 100 p.
- Federal Interagency Committee for Wetland Delineation. 1989. Federal manual for identifying and delineating jurisdictional wetlands. US Army Corps of Engineers, US Environmental Protection Agency, USDI Fish and Wildlife Service, and USDA Soil Conservation Service Cooperative Technical Publication, Washington, DC, USA. 76 p.
- Hansen, Paul L. and James B. Hall. 2002. Classification and management of USDI Bureau of Land Management's riparian and wetland sites in eastern and southern Idaho. Bitterroot Restoration, Inc., Corvallis, Montana, USA. 304 p.
- Hansen, Paul L., Robert D. Pfister, Keith Boggs, Bradley J. Cook, John Joy, and Dan K. Hinckley. 1995. Classification and management of Montana's riparian and wetland sites. Miscellaneous Publication No 54. Montana Forest and Conservation Experiment Station, School of Forestry, University of Montana, Missoula, Montana, USA. 646 p.
- Hansen, Paul L., William H. Thompson, J. Gant Massey, and Max Thompson. 2008. Classification and management of upland, riparian, and wetland sites of USDI Bureau of Land Management's Miles City Field Office, eastern Montana USA. Prepared for the USDI Bureau of Land Management Miles City Field Office, Miles City, Montana, USA. 640 p. plus 91 p. of Appendix B (Indicator Species).
- Johnson, R. R., and S. W. Carothers. 1980. Riparian habitats and recreation: interrelationships and impacts in the Rocky Mountain region. Produced under agreement 53-82 FT-0-125 of the Eisenhower Consortium for Western Environmental Forestry Research, Fort Collins, Colorado, USA. 109 p.
- Kent, Donald M. 1994. Applied wetlands science and technology. Donald M. Kent, editor. CRC Press, Inc., Lewis Publishers, Boca Raton, Florida, USA. 436 p.
- Kovalchik, Bernard L. 1987. Riparian zone associations: Deschutes, Ochoco, Fremont, and Winema National Forests. USDA Forest Service Region 6 Ecology Technical Paper 279-87. Pacific Northwest Region, Portland, Oregon, USA. 171 p.
- Mitsch, William J., and James G. Gosselink. 1993. Wetlands. Second Edition. Van Nostrand Reinhold, Publishers, New York, New York, USA. 722 p.
- Prichard, Don, Hugh Barrett, Jim Cagney, Ron Clark, Jim Fogg, Karl Gebhardt, Paul L. Hansen, Brenda Mitchell, and Dan Tippy. 1993 (revised 1995). Riparian area management: Process for assessing proper functioning condition. Technical Reference 1737-9. USDI Bureau of Land Management Service Center. Denver, Colorado. USA. 52 p.
- Prichard, Don, John Anderson, Cindy Correll, Jim Fogg, Karl Gebhardt, Russ Krapf, Steve Leonard, Brenda Mitchell, Janice Staats. 1998. Riparian area management: A user guide to assessing proper functioning condition and the supporting science for lotic areas. Technical Reference 1737-15. USDI Bureau of Land Management Service Center. Denver, Colorado. USA. 127 p.

- Reed, Porter B., Jr. 1988. National list of plant species that occur in wetlands: Northwest (Region 9). US Fish and Wildlife Service Biological Report 88 (26.9). USDI Fish and Wildlife Service, Research and Development, Washington, DC, USA. 89 p.
- Shaw, S. P., and C. G. Fredine. 1956. Wetlands of the United States: Their extent and their value for waterfowl and other wildlife. USDI Fish and Wildlife Service, Circular 39. Washington, DC, USA. 67 p.
- Stewart, R. E., and H. A. Kantrud. 1972. Classification of natural ponds and lakes in the glaciated prairie region. USDI Fish and Wildlife Service, Research Publication 92. 57 p.
- Windell, John T., Beatrice E. Willard, David J. Cooper, Susan Q. Foster, Christopher F. Knud-Hansen, Lauranne P. Rink, and George N. Kiladis. 1986. An ecological characterization of Rocky Mountain montane and subalpine wetlands. USDI Fish and Wildlife Service Biological Report 86(11). National Ecology Center, Division of Wildlife and Contaminant Research, Fish and Wildlife Service, US Department of the Interior, Washington, DC, USA. 298 p.